



Committee on Earth Observation Satellites


# GHG White Paper

David Crisp, NASA/JPL, CEOS AC-VC

CEOS AC-VC

College Park, MD, USA

2-4 May 2018





## Chapters

Contributions from participants at the CGMS and CEOS AC-VC meeting were incorporated into the White Paper structure:

### Executive Summary

- 1: Introduction
- 2: Using atmospheric GHG measurements to improve inventories
- 3: Space-based GHG measurement capabilities and near term plans
- 4: Lessons Learned from SCIAMACHY, GOSAT and OCO-2
- 5: Integrating GHG Satellites into Operational Constellations
- 6: Towards an operational constellation measuring anthropogenic CO<sub>2</sub> emissions
- 7: The Transition from Science to Operations
- 8: Conclusions



- **SCIAMACHY (2002-2012)** – First sensor to measure  $O_2$ ,  $CO_2$ , and  $CH_4$  using reflected NIR/SWIR sunlight
  - Regional-scale maps of  $X_{CO_2}$  and  $X_{CH_4}$  over continents
- **GOSAT (2009 ...)** – First Japanese GHG satellite
  - FTS optimized for high spectral resolution over broad spectral range, yielding  $CO_2$ ,  $CH_4$ , and chlorophyll fluorescence (SIF)
- **OCO-2 (2014 ...)** – First NASA satellite to measure  $O_2$  and  $CO_2$  with high sensitivity, resolution, and coverage
  - High resolution imaging grating spectrometer small ( $< 3 \text{ km}^2$ ) footprint and rapid sampling ( $10^6$  samples/day)
- **TanSat (2016 ...)** - First Chinese GHG satellite
  - Imaging grating spectrometer for  $O_2$  and  $CO_2$  bands and cloud & aerosol Imager
  - In-orbit checkout formally complete in August 2017






- **Feng Yun 3D (2017)** – Chinese GHG satellite on an operational meteorological bus
  - GAS FTS for O<sub>2</sub>, CO<sub>2</sub>, CH<sub>4</sub>, CO, N<sub>2</sub>O, H<sub>2</sub>O
- **Sentinel 5p (2017)** - Copernicus pre-operational Satellite
  - TROPOMI measures O<sub>2</sub>, CH<sub>4</sub> (1%), CO (10%), NO<sub>2</sub>, SIF
  - Imaging at 7 km x 7 km resolution, daily global coverage


---


- **Gaofen 5 (2018)** - 2<sup>nd</sup> Chinese GHG Satellite
  - Spatial heterodyne spectrometer for O<sub>2</sub>, CO<sub>2</sub>, and CH<sub>4</sub>
- **GOSAT-2 (2018)** – Japanese 2<sup>nd</sup> generation satellite
  - CO as well as CO<sub>2</sub>, CH<sub>4</sub>, with improved precision (0.125%), and active pointing to increase number of cloud free observation
- **OCO-3 (2019\*)** – NASA OCO-2 spare instrument, on ISS
  - First CO<sub>2</sub> sensor to fly in a low inclination, precessing orbit





- **CNES/UK MicroCarb (2021+)** – compact, high sensitivity
  - Imaging grating spectrometer for  $O_2 A$ ,  $O_2 \ ^1\Delta_g$ , and  $CO_2$
  - ~1/2 of the size, mass of OCO-2, with 4.5 km x 9 km footprints
- **CNES/DLR MERLIN (2021+)** - First  $CH_4$  LIDAR (IPDA)
  - Precise (1-2%)  $X_{CH_4}$  retrievals for studies of wetland emissions, inter-hemispheric gradients and continental scale annual  $CH_4$  budgets
- **NASA GeoCarb (2022\*)** – First **GEO GHG** satellite
  - Imaging spectrometer for  $X_{CO_2}$ ,  $X_{CH_4}$ ,  $X_{CO}$  and SIF
  - Stationed above  $85^\circ W$  for North/South America
- **Sentinel 5A,5B,5C (2022)** - Copernicus operational services for air quality and  $CH_4$ 
  - Daily global maps of  $X_{CO}$  and  $X_{CH_4}$  at  $< 8\text{ km} \times 8\text{ km}$





- A broad range of GHG missions will be flown over the next decade.
- Most are “science” missions, designed to identify optimal methods for measuring CO<sub>2</sub> and CH<sub>4</sub>, not “operational” missions designed to deliver policy relevant GHG products focused on anthropogenic emissions




The CO<sub>2</sub> and CH<sub>4</sub> measurement requirements in the 2011 update for the Global Climate Observing System (GCOS) Systematic Observation Requirements for Satellite-Based Data Products for Climate (GCOS, 2011) and GCOS 2016 Implementation Plan (GCOS, 2016) were adopted as targets for a future GHG constellation.

Variable / Parameter	Horizontal Resolution	Vertical Resolution	Temporal Resolution	Accuracy	Stability/Decade*	Stability/Decade**
Tropospheric CO <sub>2</sub> column	5-10km	N/A	4 h	1 ppm	0.2 ppm	1.5 ppm
Tropospheric CO <sub>2</sub>	5-10 km	5 km	4 h	1 ppm	0.2 ppm	1.5 ppm
Tropospheric CH <sub>4</sub> column	5-10 km	N/A	4 h	10 ppb	2 ppb	7 ppb
Tropospheric CH <sub>4</sub>	5-10 km	5 km	4 h	10 ppb	2 ppb	0.7 ppb
Stratospheric CH <sub>4</sub>	100-200 km	2 km	Daily	5%	0.30%	0.30%

\* from GCOS 2011

\*\* from GCOS 2016





The coverage, resolution, and precision requirements could be achieved with a constellation that incorporates

- A constellation of 3 (or more) satellites in LEO with
  - A broad ( $> 200$ ) km swath with a mean footprint size  $< 4 \text{ km}^2$
  - A single sounding random error near 0.5 ppm, and vanishing small regional scale bias ( $< 0.1 \text{ ppm}$ ) over  $> 80\%$  of the sunlit hemisphere
  - One (or more) satellites carrying ancillary sensors (CO, NO<sub>2</sub>, CO<sub>2</sub> and/or CH<sub>4</sub> Lidar)
- A constellation with 3 (or more) GEO satellites
  - Monitor diurnally varying processes (e.g. rush hours, diurnal variations in the biosphere)
  - Stationed over Europe/Africa, North/South America, and East Asia
- This constellation could be augmented with one or more HEO satellites to monitor carbon cycle changes in the high arctic



- Copernicus Sentinel CO<sub>2</sub> (2025+)
  - 3 or 4 LEO satellites in an operational GHG constellation
  - Primary instruments measure O<sub>2</sub> (0.76 μm A-band), CO<sub>2</sub> (1.61 and 2.06 μm), and NO<sub>2</sub> (0.450 μm) at a spatial resolution of 2 km x 2 km along a broad (200-300 km) swath
  - A dedicated cloud/aerosol instrument is also under consideration
- TanSat-2 Constellation
  - 6 satellites, with 3 flying in morning sun-synchronous orbits and 3 flying in afternoon sun-synchronous orbits
  - primary GHG instrument on each satellite with measure CO<sub>2</sub> (1.61 and 2.06 μm), CH<sub>4</sub> and CO (2.3 μm) as well as the O<sub>2</sub> A-band (0.76 μm) across a 100-km cross-track swath



- Because of the unprecedented requirements for precision and accuracy, the space based elements of the an operational GHG constellation architecture must be accompanied by
  - Rigorous pre-launch and on-orbit measurement calibration and product validation methods that evolve to meet emerging needs
  - Continuous refinements in remote sensing retrieval and flux inversion modeling methods that improve the products over time
- CEOS could play an essential role in coordinating these activities among its partner agencies
- Any operational architecture will also have to address
  - orbit and mission coordination, data distribution, data exchange, and data format requirements
  - Training and capacity building and public outreach will be needed to fully exploit the value of the space based GHG measurements
- CEOS should collaborate with CGMS and other operational organizations to foster the development of these capabilities



On “placing the space segment in the broader context of a fully sustained system for CO2 monitoring”

The EC proposes to organise a dedicated discussion workshop:

- Bringing together these different stakeholders to define best practices and synergies
- Exploring possibilities for common approaches to some of the system development.
- This would also require the strong engagement of CGMS as well as CEOS Associate members such as the WMO.

June 18-19<sup>th</sup> European Commission – JRC , Ispra (IT)

- **9 CEOS Agencies confirmed attendance 2-4 people each, 3 CEOS associates – around 40 attendees in total**



Monday 18<sup>th</sup> June

Tuesday 19<sup>th</sup> June

AM

## Introduction & Objectives

**International Programmes:**  
WMO, GCOS, GEO

**Ongoing Activities:**  
AC-VC Whitepaper, Inversion Modelling  
Workshop, IPCC TFI, Copernicus...

**Agencies Perspectives:**  
CNES, CSA, CMA, DLR, NOAA, NASA,  
JAXA, EU (EC, ESA, EUM), UKSA

**General & Crosscutting Aspects:**  
discussion on issues such as terminology  
System defn., needs and requirements

PM

AM

## Review of Previous Day


**Break out Groups:**  
Space-Modelling,  
Space-In-situ,  
Space-Inventory

## Reports from Breakout Groups

**Consolidation of Recommendations and Identified Best Practices**

PM

## Conclusion and Follow-up



# Overview of the Core Elements of the Copernicus CO<sub>2</sub> Emission Monitoring & Verification Support capacity

